

Prevalence of Metabolic Syndrome in a Division III Football Team

James M. Obler, Kelly A. Borgerding, Amy Olson, Ph.D., RD., LD., & Manuel Campos Ph.D.

College of Saint Benedict/Saint John's University Department of Nutrition and Biology

Introduction:

- Football players, especially linemen, strive for an increase in mass and strength to increase performance which may be associated with increased abdominal fat stores (1).
- Metabolic syndrome [MS] is a clustering of clinical symptoms including increased abdominal obesity, high systolic blood pressure [SBP], elevated triglycerides [TG], low high density lipoproteins [HDLs], and elevated fasting glucose [FBG].
- Three of these five criteria must be present in order for MS to be clinically diagnosed according to the American Heart Association and National Heart, Lung, and Blood Institute.
- MS increases the risk of cardiovascular disease and diabetes (2).

Purpose:

- The purpose of this study was to examine the prevalence of MS in a young, athletic, Division III collegiate football cohort.




Methods:

- IRB approval and informed consent was obtained for each of the 22 DIII football players (linemen, n = 15; non-linemen, n = 7).
- Subjects completed a three-day food log to record food and beverage intake.
- Height, weight [WT], waist circumference [WC], BP, FBG and lipids were measured.
- Diet records were analyzed using Diet Analysis Plus 10.0 to determine macro and micronutrient intakes.
- Individual subjects were categorized into three separate groups based upon MS risk factor [RF] prevalence (Group 1=0, 1 RF, Group 2=2 RF, Group 3=3+ RF).


Acknowledgments:

- We would like to thank head coach Gary Fasching, Richard Wielkiewicz, and the involved Saint John's football players for their participation and cooperation.


Significant MS Risk Factor ANOVA Measures



Linemen and Nonlinemen Risk Factor Prevalence



Team and NHANES Risk Factor Prevalence



Results:

- Prevalence of MS within this cohort was 23% (5 of 22); 27% (4 of 15) of linemen and 14% of non-linemen (1 of 7).
- Pearson correlation coefficients and an analysis of variance were used for statistical analyses.
- Significant ANOVA variations between groups include WT ($p = .002$), WC ($p = .002$), SBP ($p = .015$), and TG ($p = .005$).
- Linemen compared to non-linemen met MS criteria for WC [73% vs. 0%], HDL [40% vs. 14%], and SBP [80% vs. 57%].
- WT significantly correlated with WC ($r = .898$ $p = .000$) and SBP ($r = .494$ $p = .019$). SBP significantly correlated with dietary total fat ($r = .439$ $p = .041$) and saturated fat intake ($r = .427$ $p = .047$).
- SBP for all subjects averaged 40% higher than an age matched cohort from NHANES.



Conclusions:

- Despite being physically active, the prevalence of MS amongst linemen was 27% and 14% among non-linemen.
- Linemen were larger and had more MS risk factors within this cohort.
- These results may predict future health problems in DIII football players since the MS prevalence in NFL linemen retirees is ~60% (3).
- Nutritional counseling may help reduce cardiometabolic risk factors given their WT and WC, and the significant correlation with dietary total and saturated fat and SBP.
- Future research should examine whether the presence of MS risk factors also affects performance.

References:

- Wilkerson, et al. (2010). *Journal of Athletic Training*, 45, 67-74.
- Buell, et al. (2008). *Journal of Athletic Training*, 43, 608-16.
- Miller, et al. (2008). *American Journal of Cardiology*, 101, 1281-4.